

=> FILE REG

FILE 'REGISTRY' ENTERED AT 20:31:21 ON 15 DEC 2010
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2010 American Chemical Society (ACS)

=> D HIS

FILE 'LREGISTRY' ENTERED AT 20:13:44 ON 15 DEC 2010
L1 STR

FILE 'REGISTRY' ENTERED AT 20:20:12 ON 15 DEC 2010
L2 SCR 2043
L3 0 S L1 AND L2

FILE 'LREGISTRY' ENTERED AT 20:20:22 ON 15 DEC 2010
L4 STR L1


FILE 'REGISTRY' ENTERED AT 20:22:17 ON 15 DEC 2010
L5 3 S L4 AND L2
L6 STR L4
L7 0 S L6 AND L2
L8 50 S L4
L9 1187 S L4 FUL
SAV L9 CON424/A
E PMS/CI
L10 1371382 S E3
L11 35 S L10 AND L9

FILE 'HCA' ENTERED AT 20:29:05 ON 15 DEC 2010
L12 14 S L11
L13 10 S 1802-2003/PY,PRY,AY AND L12

FILE 'REGISTRY' ENTERED AT 20:31:21 ON 15 DEC 2010

=> D L9 QUE STAT

L4 STR



VAR G1=O/S

NODE ATTRIBUTES:

DEFAULT MLEVEL IS ATOM

DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:

RING(S) ARE ISOLATED OR EMBEDDED

NUMBER OF NODES IS 12

STEREO ATTRIBUTES: NONE

L9 1187 SEA FILE=REGISTRY SSS FUL L4

100.0% PROCESSED 1482003 ITERATIONS (8 INCOMPLETE) 1187 ANSWERS
SEARCH TIME: 00.00.02

=> FILE HCA

FILE 'HCA' ENTERED AT 20:31:33 ON 15 DEC 2010

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2010 AMERICAN CHEMICAL SOCIETY (ACS)

=> D L13 1-10 BIB ABS HITSTR HITRN RE

L13 ANSWER 1 OF 10 HCA COPYRIGHT 2010 ACS on STN

AN 141:382155 HCA Full-text

TI Electrode active material for electrochemical device

IN Inatomi, Yuu; Shimada, Mikinari

PA Matsushita Electric Industrial Co., Ltd., Japan

SO U.S. Pat. Appl. Publ., 12 pp.

CODEN: USXXCO

DT Patent

LA English


FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	US 20040214082	A1	20041028	US 2004-827424	20040420
	EP 1478040	A1	20041117	EP 2004-252356	20040421
	EP 1478040	B1	20080423		
	JP 2004342605	A	20041202	JP 2004-124954	20040421
	JP 4468058	B2	20100526		
	CN 1540786	A	20041027	CN 2004-10035314	20040422
	CN 1264237	C	20060712		
PRAI	JP 2003-116843	A	20030422		

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OS MARPAT 141:382155

GI



AB An electrochem. device of the present invention comprises a pos. electrode, a neg. electrode and an electrolyte, wherein at least one of the pos. electrode and the neg. electrode includes a compd. having a structure represented by the general formula (I); where X is a S atom, a N atom, or an O atom, each of R1 to R4 is independently a linear or cyclic aliph. group, a H atom, a -OH group, a cyano group, an amino group, a nitro group, or a nitroso group; each of R5 and R6 is independently a linear or cyclic aliph. group, the aliph. group includes ≥ 1 selected from the group consisting of an O atom, a N atom, a S atom, a Si atom, a P atom, a B atom, and a halogen atom. It is thereby possible to obtain a lightwt. and high energy-d. electrochem. device having an excellent cycle characteristic.

IT 782452-24-8P 782452-26-0P

(electrode active material for electrochem. device)


RN 782452-24-8 HCA

CN 1,3-Dithiole, 2,2'-(1,2-ethanediyldiene)bis-, polymer with ethene
(9CI) (CA INDEX NAME)

CM 1

CRN 88312-71-4

CMF C8 H6 S4



CM 2

CRN 74-85-1

CMF C2 H4




RN 782452-26-0 HCA

CN 1,3-Dithiole, 2-(1,3-dithiol-2-ylideneethylidene)-4-(2-propynyl)-, polymer with ethyne (9CI) (CA INDEX NAME)

CM 1

CRN 782452-25-9


CMF C11 H8 S4



CM 2

CRN 74-86-2

CMF C2 H2



IT 782452-24-8P 782452-26-0P

(electrode active material for electrochem. device)

OSC.G 3 THERE ARE 3 CAPLUS RECORDS THAT CITE THIS RECORD (8 CITINGS)

L13 ANSWER 2 OF 10 HCA COPYRIGHT 2010 ACS on STN

AN 137:370466 HCA Full-text

TI Acrylic esters containing 1,3-dithiolane linkage and their use for optical materials

IN Nakamura, Mitsuo; Imai, Masao; Otsuji, Atsuo

PA Mitsui Chemicals, Inc., Japan

SO PCT Int. Appl., 104 pp.

CODEN: PIXXD2

DT Patent

LA Japanese

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	WO 2002092591	A1	20021121	WO 2002-JP4695	20020515
	AU 2002308979	A1	20021125	AU 2002-308979	20020515
	AU 2002308979	B2	20040923		
	JP 2003160615	A	20030603	JP 2002-140663	20020515
	JP 3820183	B2	20060913		
	EP 1388540	A1	20040211	EP 2002-769605	20020515
	EP 1388540	B1	20070905		
	CN 100344623	C	20071024	CN 2002-801665	20020515
	US 20040102592	A1	20040527	US 2003-332742	20030113
	US 6835844	B2	20041228		
PRAI	JP 2001-144539	A	20010515		
	JP 2001-144540	A	20010515		
	JP 2001-266422	A	20010903		
	JP 2001-281859	A	20010917		
	WO 2002-JP4695	W	20020515		

OS MARPAT 137:370466

AB Acrylic esters which are polymerizable in a short time and useful for optical resins with good heat, mech. and refractive properties are diacrylic esters having dithiolane linkage. Acid condensation of 2,3-dimercapto-1-propanol with glyoxal, treating the resulting 2,2'-bis[(4-hydroxymethyl)dithiolane] with thiourea, hydrolysis of the thiouronium salt, treating the resulting dithiol with 3-chloropropionic acid, and treating with Et3N gave 2,2'-bis[(4-acryloylthiomethyl)dithiolane]. The dithiolane was UV cured with Darocur 1173 in a cell, giving a transparent test piece with refractive index 1.681, Abbe no. 34.6, sp. gr. 1.4, and glass temp. 123°.

IT ~~475577-82-3P~~

(manuf. of acrylic esters contg. 1,3-dithiolane linkage for optical materials)


RN 475577-82-3 HCA

CN 2-Propenethioic acid, S,S'-[1,2-ethanediytlebis(1,3-dithiolane-2,4-diylmethylen)] ester, homopolymer (9CI) (CA INDEX NAME)

CM 1

CRN 475577-77-6

CMF C16 H22 O2 S6



IT 475577-82-3P

(manuf. of acrylic esters contg. 1,3-dithiolane linkage for optical materials)

RE CITED REFERENCES

- (1) Essilor Int Cie Gen Optique; JP 2000509075 A 1998
- (2) Essilor Int Cie Gen Optique; US 20020049289 A 1998 HCA
- (3) Essilor Int Cie Gen Optique; FR 2759369 A 1998 HCA
- (4) Essilor Int Cie Gen Optique; EP 901486 A1 1998 HCA
- (5) Essilor Int Cie Gen Optique; WO 9835955 A1 1998 HCA
- (6) Essilor Int Cie Gen Optique; AU 9866269 A 1998 HCA
- (7) Hoya Corp; JP 03-215801 A 1991 HCA
- (8) Mitsui Chem Inc; WO 0192252 A1 2000
- (9) Mitsui Chem Inc; EP 1057808 A2 2000 HCA
- (10) Mitsui Chem Inc; CN 1283614 A 2000 HCA
- (11) Mitsui Chem Inc; JP 2001316355 A 2000 HCA
- (12) Mitsui Chem Inc; JP 200148877 A 2000
- (13) Mitsui Petrochem Ind Co Ltd; JP 04-321662 A 1992 HCA

OSC.G 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (7 CITINGS)

L13 ANSWER 3 OF 10 HCA COPYRIGHT 2010 ACS on STN

AN 137:140883 HCA Full-text

TI Poly(dibenzylidenetetrathiapentalene): A Redox-Active, Linearly Extended TTF Polymer

AU Divisia-Blohorn, Bernadette; Genoud, Francoise; Salhi, Fouad; Mueller, Harald

CS Laboratoire d'Electrochimie Moleculaire et Structure des Interfaces and Laboratoire de Physique des Metaux Synthetiques, Unite Mixte de Recherche CEA-CNRS-Univ.-J. Fourier N° 5819, Grenoble, 38054, Fr.

SO Journal of Physical Chemistry B (2002), 106(26), 6646-6651
CODEN: JPCBFK; ISSN: 1089-5647

PB American Chemical Society

DT Journal

LA English

AB The π -conjugated polymers composed of vinylogous tetrathiafulvalene (TTF) units were obtained by electrochem. polymn. of 2,5-di(benzylidene)-1,3,4,6-tetrathiapentalene or of its corresponding dimer 1,2-bis(5-benzylidene[1,3]-dithiolo[4,5-d]1,3-dithiole-2-ylidene)-1,2-diphenylethane. The polymers were characterized by cyclic voltammetry (CV), in situ ESR spectroscopy and in situ cond. measurements to gain insight into the charge-discharge behavior and the electronic properties as a function of applied potential. Our results suggest that both compds. behave as conducting polymers exhibiting multistep redox activity. In situ ESR studies show that charge-

discharge phenomena differ in the nature of the charge storage configuration depending on whether they are investigated during a cathodic or anodic scan. Redn. of a polymer film generated at an applied potential of +1.6 V vs SCE allows for the detection of three clearly discernible maxima of the spin d. at 0.60, 0.95, and about 1.25 V vs Ag, while subsequent reoxidn. gives rise to only one spin d. max. at around 0.65 V vs Ag. Coulometric studies suggest that the spin d. max. at around 0.60-0.65 V vs Ag is assocd. with one pos. charge per tetrathiapentalene unit of poly-1 while each monomer bears up to two charges at a potential of 1.6 V vs Ag. In situ and conventional four-probe measurements of the elec. cond. gave values of around $5 \times 10^{-5} \text{ S cm}^{-1}$.

IT 444874-87-7P

(prepn. and electrochem. properties of)


RN 444874-87-7 HCA

CN [1,3]Dithiolo[4,5-d]-1,3-dithiole,
2,2'-(1,2-diphenyl-1,2-ethanediylidene)bis[5-(phenylmethylene)-,
homopolymer (9CI) (CA INDEX NAME)

CM 1

CRN 189077-89-2

CMF C36 H22 S8



IT 444874-87-7P

(prepn. and electrochem. properties of)

RE CITED REFERENCES

- (1) Abruna, H; Coord Chem Rev 1988, V86, P135 HCA
- (2) Abruna, H; Electroresponsive Molecular and Polymeric Systems, Chapter 3 1988, V1
- (3) Adam, M; Adv Mater 1994, V6, P439
- (4) Adamo, C; J Phys Chem B 1999, V103, P6863 HCA
- (5) Aubert, P; J Electroanal Chem 1999, V470, P77 HCA
- (6) Becher, J; Electronic Materials: The Oligomer Approach, Chapter 2.2 1998
- (7) Chidsey, C; J Phys Chem 1986, V90, P1484
- (8) Feldman, B; Adv Mater 1995, V7, P48
- (9) Frenzel, S; J Mater Chem 1995, V5, P1529 HCA
- (10) Froyer, G; Synth Met 1989, V33, P381 HCA
- (11) Genoud, F; J Chim Phys 1990, V87, P57 HCA
- (12) Genoud, F; Phys Rev Lett 1985, V55, P118 HCA
- (13) Graf, D; J Am Chem Soc 1999, V119, P5888
- (14) Gruber, H; Synth Met 1991, V41-43, P2331
- (15) Gruber, H; Synth Met 1991, V44, P55 HCA
- (16) Hapiot, P; J Phys Chem A 1999, V103, P11221 HCA
- (17) Hapiot, P; Synth Met 1999, V101, P77 CAPLUS


- (18) Hill, M; Chem Mater 1992, V4, P1106 HCA
- (19) Inzelt, G; J Electroanal Chem Interfacial Electrochem 1983, V159, P443
HCA
- (20) Jestin, I; J Am Chem Soc 1998, V120, P8150 HCA
- (21) Kaufman, F; J Am Chem Soc 1979, V102, P483
- (22) Kossmehl, G; Macromolecules 1982, V9, P541
- (23) Lane, R; J Phys Chem 1973, V77, P1401 HCA
- (24) Laviron, E; J Electroanal Chem 1972, V39, P1 HCA
- (25) Laviron, E; J Electroanal Chem 1974, V52, P395 HCA
- (26) Laviron, E; J Electroanal Chem 1979, V105, P25 HCA
- (27) Levi, M; J Phys Chem B 1999, V103, P1499 HCA
- (28) Levillain, E; J Am Chem Soc 1999, V121, P8760 HCA
- (29) Miller, L; Acc Chem Res 1996, V29, P417 HCA
- (30) Muller, H; Chem Commun 1999, P1407 HCA
- (31) Muller, H; J Chim Phys 1998, V95, P1184 HCA
- (32) Muller, H; Synth Met 1997, V84, P445
- (33) Muller, H; Synth Met 1999, V101, P75
- (34) Muller, H; Synth Met 2001, V119, P195 HCA
- (35) Muller, H; Tetrahedron Lett 1997, V38, P3215 HCA
- (36) Murray, R; J Am Chem Soc 1985, V107, P872
- (37) Nechtschein, M; Handbook of Conducting Polymers, 2nd ed, Chapter 9
1998
- (38) Neudeck, A; J Phys Chem B 1999, V103, P912 HCA
- (39) Neudeck, A; Synth Met 1999, V107, P143 HCA
- (40) Ofer, D; J Am Chem Soc 1990, V112, P7869 HCA
- (41) Park, S; Handbook of Organic Conductive Molecules and Polymers,
Chapter 9 1997, V3
- (42) Pham, M; J Electrochem Soc 1993, V140, P912
- (43) Pickup, P; J Mater Chem 1999, V9, P1641 HCA
- (44) Pittman, C; Macromolecules 1976, V9, P360 HCA
- (45) Pittman, C; Macromolecules 1979, V12, P541 HCA
- (46) Pullen, A; Macromolecules 2001, V34, P812 HCA
- (47) Rapta, P; J Chem Soc, Faraday Trans 1998, V94, P3625 HCA
- (48) Roncali, J; J Mater Chem 1997, V7, P2307 HCA
- (49) Roth, H; Prog Colloid Polym Sci 1988, V78, P75 HCA
- (50) Roth, H; Prog Colloid Polym Sci 1989, V80, P254 HCA
- (51) Rulkens, R; J Am Chem Soc 1996, V118, P12683 HCA
- (52) Salhi, F; Ph D Thesis, Universite Joseph Fourier 1999
- (53) Schukat, G; Sulfur Rep 1993, V14, P245 HCA
- (54) Shimizu, T; Synthesis 1998, V3, P259
- (55) Thobie-Gauthier, C; Macromolecules 1993, V26, P4094
- (56) Trinh, V; J Prakt Chem 1989, V331, P826 HCA
- (57) Ueno, Y; Chem Lett 1975, P603 HCA
- (58) Yamamoto, T; J Mater Chem 1997, V7, P1967 HCA
- (59) Yamamoto, T; J Mater Chem 1997, V7, P1967 HCA
- (60) Yamamoto, T; Macromolecules 1997, V30, P5390 HCA
- (61) Zhou, M; J Phys Chem B 1999, V103, P8443 HCA
- (62) Zhou, M; J Phys Chem B 1999, V103, P8451 HCA
- (63) Zotti, G; Chem Mater 1991, V3, P62 HCA
- (64) Zotti, G; Chem Mater 2000, V12, P2996 HCA

OSC.G 3 THERE ARE 3 CAPLUS RECORDS THAT CITE THIS RECORD (3 CITINGS)

L13 ANSWER 4 OF 10 HCA COPYRIGHT 2010 ACS on STN
 AN 136:200988 HCA Full-text
 TI 4-Methylene-1,3-dioxolanes as odorless crosslinking agents
 IN Hartl, Helmut; Frings, Rainer B.; Grahe, Gerwald F.
 PA Dainippon Ink and Chemicals, Inc., Japan
 SO Eur. Pat. Appl., 16 pp.
 CODEN: EPXXDW
 DT Patent
 LA English
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	EP 1182201	A1	20020227	EP 2000-118116	20000825
	EP 1182201	B1	20041124		
	ES 2233255	T3	20050616	ES 2000-118116	20000825
	CA 2355749	A1	20020225	CA 2001-2355749	20010821
	CA 2355749	C	20080923		
	US 20020045674	A1	20020418	US 2001-934655	20010823
	US 6844375	B2	20050118		
	JP 2002155073	A	20020528	JP 2001-254419	20010824
PRAI	EP 2000-118116	A	20000825		

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT
 OS MARPAT 136:200988
 GI



AB The invention relates to 4-methylene-1,3-dioxolanes of the general formula I wherein R1 denotes H, C5-6-cycloalkyl or C1-4-alkyl; m and n, which may be the same or different, denote 0 or 1, whereby $m \leq n$, o denotes 2, 3 or 4 depending on the valency of the group X; and X denotes a C-C single bond, straight-chain or branched C1-18-alkylene, C5-6-cycloalkylene, C8-18-arylalkylene, $-\text{CH}_2(\text{OCH}_2\text{CH}_2)_p\text{OCH}_2-$, $-\text{CH}_2(\text{OCH}(\text{CH}_3)\text{CH}_2)_p\text{OCH}_2-$, wherein p is an integer from 0 to 100, a process for their prodn. and intermediate products used. Moreover, compns. capable of emission-free, photocationic crosslinking, which comprise 4-methylene-1,3-dioxolanes of the general formula I and their use for the prodn. of solvent-resistant and transparent films. Thus, heating 500 g a .apprx.50% aq. glutardialdehyde soln. with 600 g 3-chloro-1,2-propanediol and .apprx.400 mL PhMe to boiling while removing water (340 mL) over 3-4 h, cooling to room temp., washing, drying, removing PhMe by a rotary evaporator and distg. the residue gave 627 g (88%) 1,3-bis(4-chloromethyl-1,3-dioxolane-2-yl)propane (II) with b.p. 138-140° (5x10-3 mbar). Adding II to K tert-butyrate dissolved in dry THF while

maintaining at below 50° then stirring at room temp. for 24 h and working up gave

1,3-bis(4-methylene-1,3-dioxolane-2-yl)propane.

IT ~~401794-52-3P 401794-55-6P 401794-63-6P~~

(manuf. of 4-methylene-1,3-dioxolanes as odorless crosslinking agents)


RN 401794-52-3 HCA

CN 1,3-Dioxolane, 2,2'-(1,2-ethanediyl)bis[2-methyl-4-methylene-, homopolymer (9CI) (CA INDEX NAME)

CM 1

CRN 401794-51-2

CMF C12 H18 O4




RN 401794-55-6 HCA

CN 1,3-Dioxolane, 2,2'-(1,3-propanediyl)bis[4-methylene-, polymer with 2,2'-(1,2-ethanediyl)bis[2-methyl-4-methylene-1,3-dioxolane] (9CI) (CA INDEX NAME)

CM 1

CRN 401794-51-2


CMF C12 H18 O4



CM 2

CRN 401793-74-6

CMF C11 H16 O4




RN 401794-63-6 HCA

CN 1,3-Dioxolane, 2,2'-(1,2-ethanediyl)bis[2-methyl-4-methylene-, polymer with 2,2'-[(1-methylethylidene)bis(4,1-phenyleneoxymethylene)]bis[4-methylene-1,3-dioxolane] (9CI) (CA INDEX NAME)

CM 1

CRN 401794-51-2


CMF C12 H18 O4



CM 2

CRN 401793-97-3

CMF C25 H28 O6




IT 401794-52-3P 401794-55-6P 401794-63-6P

(manuf. of 4-methylene-1,3-dioxolanes as odorless crosslinking agents)

RE CITED REFERENCES

- (1) Abdullaev, N; 1977 HCA
- (2) Abdullaev, N; KHIM PRIR SOEDIN 1976, V6, P828
- (3) Dynamit Nobel; GB 724032 A
- (4) Sokolov, G; 1965
- (5) Sokolov, G; 1972 HCA
- (6) Sokolov, G; KHIM-FARM ZH 1972, V6(2), P10 HCA
- (7) Sokolov, G; LATVIJAS PSR ZINATNU AKAD VESTIS, KIM SER 1964, V6, P667

L13 ANSWER 5 OF 10 HCA COPYRIGHT 2010 ACS on STN
 AN 133:80761 HCA Full-text
 TI New electrochemically synthesized copolymers:
 poly(difluorenyl-ethylenes)
 AU Lorcy, Dominique; Rault-Berthelot, Joelle; Poriel, Cyril
 CS Synthese et Electrosynthese Organiques, UMR CNRS 6510, Universite de
 Rennes 1, Rennes, 35042, Fr.
 SO Electrochemistry Communications (2000), 2(6), 382-385
 CODEN: ECCMF9; ISSN: 1388-2481
 PB Elsevier Science B.V.
 DT Journal
 LA English
 AB Synthesis and cyclic voltammetry data of novel copolymers contg.
 difluorenyl-ethylene units in the conjugated backbone are reported. The
 dimer of 2-(dithiafulvenyl)fluorene is chem. and electrochem. formed and
 electropolymd. and voltammetry of the polymer studied. 1,2-Difluorenyl-1,2-
 dimethylethene is also electrochem. polymd.
 IT 279668-55-2P
 (electrochem. prepn. and cyclic voltammetry in CH2Cl2)
 RN 279668-55-2 HCA
 CN 1,3-Dithiole, 2,2'-(1,2-di-9H-fluoren-2-yl-1,2-ethanediylidene)bis[4,5-
 dimethyl-, homopolymer (9CI) (CA INDEX NAME)
 CM 1
 CRN 279668-52-9
 CMF C38 H30 S4



IT 279668-55-2P
 (electrochem. prepn. and cyclic voltammetry in CH2Cl2)
 RE CITED REFERENCES
 (1) Bellec, N; J Electroanal Chem 1999, V462, P137 HCA
 (2) Benahmed-Gasmi, A; Tetrahedron Lett 1995, V36, P2983 HCA
 (3) Burroughes, J; Nature 1990, V347, P539 HCA
 (4) Cho, H; Adv Mater 1997, V9, P326 HCA
 (5) de Leeuw, D; Phys World 1999, V12, P31 HCA
 (6) Friend, R; Nature 1999, V397, P121 HCA
 (7) Gonzalez, S; J Org Chem 1999, V64, P3498 HCA

(8) Hapiot, P; J Phys Chem 1996, V100, P14823 HCA
 (9) Hascoat, P; J Org Chem 1997, V62, P6086 HCA
 (10) Kido, J; Phys World 1999, V12, P27 HCA
 (11) Kraft, A; Angew Chem Int Ed Engl 1998, V37, P402
 (12) Lorcy, D; J Org Chem 1995, V60, P2443 HCA
 (13) Lorcy, D; Synth Met 1997, V86, P1831 HCA
 (14) Mayer, R; J Prakt Chem 1974, V316, P907 HCA
 (15) Minabe, M; J Org Chem 1976, V41, P1935 HCA
 (16) Moore, A; J Chem Soc Perkin Trans 1 1997, P3443 HCA
 (17) Muller, H; Tetrahedron Lett 1997, V38, P3215 HCA
 (18) Ohmori, Y; Jpn J Appl Phys 1991, V30, PL1941
 (19) Ohta, A; J Chem Soc Chem Commun 1995, P1761 HCA
 (20) Pei, Q; J Am Chem Soc 1996, V118, P7416 HCA
 (21) Ranger, M; Macromolecules 1997, V30, P7686 HCA
 (22) Rault-Berthelot, J; Electrochim Acta 1988, V33, P811 HCA
 (23) Rault-Berthelot, J; Recent Res Dev Macromol Res 1998, V3, P425 HCA
 (24) Schoberl, U; Adv Mater 1992, V4, P41
 (25) Yamashita, Y; Chem Commun 1998, P1657 HCA
 (26) Yu, W; Chem Commun 1999, P1837 HCA
 OSC.G 5 THERE ARE 5 CAPLUS RECORDS THAT CITE THIS RECORD (5 CITINGS)

L13 ANSWER 6 OF 10 HCA COPYRIGHT 2010 ACS on STN
 AN 131:18963 HCA Full-text
 TI Synthesis, X-ray Structure, and Electrochemical Oxidative Coupling
 Reactions of 1,5- and 2,6-Bis(1,4-dithiafulven-6-yl)naphthalenes
 AU Gonzalez, Susana; Martin, Nazario; Sanchez, Luis; Segura, Jose L.;
 Seoane, Carlos; Fonseca, Isabel; Cano, Felix H.; Sedo, Josep;
 Vidal-Gancedo, Jose; Rovira, Concepcio
 CS Departamento de Quimica Organica Facultad de Quimica, Universidad
 Complutense, Madrid, E-28040, Spain
 SO Journal of Organic Chemistry (~~1999~~), 64(10), 3498-3506
 CODEN: JOCEAH; ISSN: 0022-3263
 PB American Chemical Society
 DT Journal
 LA English
 OS CASREACT 131:18963
 AB Novel π -extended tetrathiafulvalene (TTF) derivs. I and II [R = H, SMe; RR =
 SCH₂CH₂S; R1 = Hexyl, Me], in which the two 1,3-dithiole units are connected
 through a naphthalene spacer, have been prepd. in high yields by Wittig-
 Horner olefination reaction of dialkoxy diformylnaphthalenes and substituted
 phosphonate esters. The electrochem. study revealed a similar behavior for
 the novel electron donor mols. I and II regardless of the position of the
 1,3-dithiole rings on the naphthalene core. The extended donors undergo an
 efficient electrooxidn. process affording new oligomeric extended TTF
 species which exhibit lower oxidn. potential values than their precursor
 donors. EPR expts. confirm the presence of the cation radical derived from
 the oligomeric TTF vinylogues and support an ECE (electrochem.-chem.-
 electrochem.) process. The structural study has been carried out by X-ray
 anal. of I [R = H; R1 = Hexyl] and semiempirical PM3 calcns. and reveals a
 distorted geometry from planarity with the naphthalene moiety forming an
 angle of .apprx.35° with the thiafulvalene rings. A good agreement was

found between the exptl. and calcd. values, thus validating the PM3 method. The chem. oxidn. of I and II with strong electron acceptors give charge transfer complexes (CTC) which were characterized by UV-vis, FTIR, and EPR spectroscopy.

IT 226698-09-5P 226698-11-9P 226698-13-1P
 226698-14-2P 226698-15-3P 226698-17-5P
 226698-18-6P 226698-19-7P 226698-20-0P
 226698-21-1P

(charge transfer complex; mol. structure and electrochem. oxidative coupling of tetrathiafulvenes contg. naphthalene spacers prepd. by Wittig-Horner olefination of dithiolyolphosphonate esters with diformylnaphthalenes)

RN 226698-09-5 HCA


CN Propanedinitrile, 2,2'-(2,5-cyclohexadiene-1,4-diylidene)bis-, compd. with poly[(1,5-dimethoxy-2,6-naphthalenediyl)[1,2-bis[4,5-bis(methylthio)-1,3-dithiol-2-yl]-1,2-ethenediyl]] (9CI) (CA INDEX NAME)

CM 1

CRN 226698-08-4

CMF (C24 H24 O2 S8)n


CCI PMS



CM 2

CRN 1518-16-7


CMF C12 H4 N4



RN 226698-11-9 HCA
 CN Propanedinitrile, 2,2'-(2,5-cyclohexadiene-1,4-diylidene)bis-, compd.
 with poly[(4,8-dimethoxy-1,5-naphthalenediyl)[1,2-bis(1,3-dithiol-2-
 yl)-1,2-ethenediyl]] (9CI) (CA INDEX NAME)


CM 1

CRN 226698-10-8
 CMF (C20 H16 O2 S4)n
 CCI PMS



CM 2

CRN 1518-16-7
 CMF C12 H4 N4




RN 226698-13-1 HCA
 CN 1,4-Cyclohexadiene-1,2-dicarbonitrile, 4,5-dichloro-3,6-dioxo-, compd.
 with poly[(1,5-dimethoxy-2,6-naphthalenediyl)[1,2-bis(1,3-dithiol-2-
 yl)-1,2-ethenediyl]] (9CI) (CA INDEX NAME)

CM 1

CRN 226698-12-0

CMF (C20 H16 O2 S4)n


CCI PMS



CM 2

CRN 84-58-2

CMF C8 C12 N2 O2



RN 226698-14-2 HCA


CN 1,4-Cyclohexadiene-1,2-dicarbonitrile, 4,5-dichloro-3,6-dioxo-, compd. with poly[(1,5-dimethoxy-2,6-naphthalenediyl)[1,2-bis[4,5-bis(methylthio)-1,3-dithiol-2-yl]-1,2-ethenediyl]] (9CI) (CA INDEX NAME)

CM 1

CRN 226698-08-4

CMF (C24 H24 O2 S8)n


CCI PMS



CM 2

CRN 84-58-2

CMF C8 C12 N2 O2



RN 226698-15-3 HCA


CN 1,4-Cyclohexadiene-1,2-dicarbonitrile, 4,5-dichloro-3,6-dioxo-, compd. with poly[(4,8-dimethoxy-1,5-naphthalenediyl)[1,2-bis(1,3-dithiol-2-yl)-1,2-ethenediyl]] (9CI) (CA INDEX NAME)

CM 1

CRN 226698-10-8

CMF (C20 H16 O2 S4)n


CCI PMS



CM 2

CRN 84-58-2

CMF C8 C12 N2 O2



RN 226698-17-5 HCA


CN 1,4-Cyclohexadiene-1,2-dicarbonitrile, 4,5-dichloro-3,6-dioxo-, compd. with poly[(4,8-dimethoxy-1,5-naphthalenediyl)[1,2-bis[4,5-bis(methylthio)-1,3-dithiol-2-yl]-1,2-ethenediyl]] (9CI) (CA INDEX NAME)

CM 1

CRN 226698-16-4

CMF (C24 H24 O2 S8)n


CCI PMS



CM 2

CRN 84-58-2

CMF C8 C12 N2 O2



RN 226698-18-6 HCA


CN Cyanamide, (2,3-dichloro-5,6-dimethyl-2,5-cyclohexadiene-1,4-diyldiene)bis-, compd. with poly[(1,5-dimethoxy-2,6-naphthalenediyl)[1,2-bis(1,3-dithiol-2-yl)-1,2-ethenediyl]] (9CI) (CA INDEX NAME)

CM 1

CRN 226698-12-0

CMF (C20 H16 O2 S4)n


CCI PMS



CM 2

CRN 157788-94-8

CMF C10 H6 C12 N4



RN 226698-19-7 HCA


CN Cyanamide, (2,3-dichloro-5,6-dimethyl-2,5-cyclohexadiene-1,4-diylidene)bis-, compd. with poly[(1,5-dimethoxy-2,6-naphthalenediyl)[1,2-bis[4,5-bis(methylthio)-1,3-dithiol-2-yl]-1,2-ethenediyl]] (9CI) (CA INDEX NAME)

CM 1

CRN 226698-08-4

CMF (C24 H24 O2 S8)n


CCI PMS



CM 2

CRN 157788-94-8

CMF C10 H6 Cl2 N4



RN 226698-20-0 HCA


CN Cyanamide, (2,3-dichloro-5,6-dimethyl-2,5-cyclohexadiene-1,4-diylidene)bis-, compd. with poly[(4,8-dimethoxy-1,5-naphthalenediyl)[1,2-bis(1,3-dithiol-2-yl)-1,2-ethenediyl]] (9CI) (CA INDEX NAME)

CM 1

CRN 226698-10-8

CMF (C20 H16 O2 S4)_n


CCI PMS



CM 2

CRN 157788-94-8

CMF C10 H6 Cl2 N4




RN 226698-21-1 HCA
 CN Cyanamide, (2,3-dichloro-5,6-dimethyl-2,5-cyclohexadiene-1,4-diylidene)bis-, compd. with poly[(4,8-dimethoxy-1,5-naphthalenediyl)[1,2-bis[4,5-bis(methylthio)-1,3-dithiol-2-yl]-1,2-ethenediyl]] (9CI) (CA INDEX NAME)

CM 1

CRN 226698-16-4

CMF (C24 H24 O2 S8)n


CCI PMS



CM 2

CRN 157788-94-8

CMF C10 H6 C12 N4



IT 226698-09-5P 226698-11-9P 226698-13-1P
226698-14-2P 226698-15-3P 226698-17-5P
226698-18-6P 226698-19-7P 226698-20-0P
226698-21-1P

(charge transfer complex; mol. structure and electrochem. oxidative coupling of tetrathiafulvenes contg. naphthalene spacers prepd. by Wittig-Horner olefination of dithiolylphosphonate esters with diformyl naphthalenes)

RE CITED REFERENCES

- (1) Akiba, K; Bull Chem Soc Jpn 1978, V51, P2674 HCA
- (2) Altomate, A; Dipartimento Geomineralogico 1992
- (3) Anon; International Tables for X-Ray Crystallography 1974, V4
- (4) Awaji, H; Chem Mater 1989, V1, P535
- (5) Benahmed-Gasmi, A; Tetrahedron Lett 1995, V36, P2983 HCA
- (6) Brisset, H; J Chem Soc Chem Commun 1994, P1765 HCA
- (7) Bryce, M; Angew Chem Int Ed Engl 1990, V29, P1450
- (8) Bryce, M; Chem Mater 1996, V8, P1182 HCA
- (9) Bryce, M; J Mater Chem 1995, V5, P1481 HCA
- (10) Bun-Hoy, N; J Org Chem 1955, V20, P1191
- (11) De Lucas, A; Tetrahedron 1998, V54, P4655 HCA
- (12) Faldt, A; Synth Met 1998, V94, P307 HCA
- (13) Fisher, P; J Am Chem Soc 1963, V85, P2694
- (14) Fourmigue, M; J Am Chem Soc 1993, V115, P3752 HCA
- (15) Garin, J; Tetrahedron Lett 1998, V39, P3577 HCA
- (16) Gonzalez, M; Tetrahedron Lett 1998, V39, P3269 HCA
- (17) Hall, S; XTAL3.2 System 1994
- (18) Hapiot, P; J Phys Chem 1996, V100, P14823 HCA
- (19) Herranz, M; Tetrahedron 1998, V54, P11651 HCA
- (20) Hunig, S; Int J Sulfur Chem C 1971, V6, P109
- (21) Iianack, M; Adv Mater 1996, V8, P663
- (22) Iiascoat, P; J Org Chem 1997, V62, P6086
- (23) Jen, A; J Chem Soc Chem Commun 1994, P2057 HCA
- (24) Khodorkovsky, V; Organic Conductors: Fundamentals and Applications 1994, P75 HCA
- (25) Lorcy, D; J Org Chem 1995, V60, P2443 HCA
- (26) Martin, N; J Org Chem 1998, V63, P1268 HCA
- (27) Martin, N; Synth Met 1994, V64, P83 HCA
- (28) Martin, N; Synth Met 1996, V76, P137
- (29) Martinez-Ripoll, M; PESOS A Computer Program for the Automatic Treatment of Weighting Schemes 1975
- (30) Miller, J; J Am Chem Soc 1986, V108, P4459 HCA
- (31) Moore, A; J Chem Soc Perkin Trans 1 1991, P157 HCA

(32) Moore, A; Synthesis 1991, P26 HCA
 (33) Nardelli, M; Comput Chem 1973, V7, P95
 (34) Ohta, A; Heterocycles 1995, V40, P123 HCA
 (35) Papavassiliou, G; Organic Conductive Molecules and Polymers 1997, P151
 (36) Roncali, J; J Mater Chem 1997, V7, P2307 HCA
 (37) Schoberl, H; Adv Mater 1992, V4, P41
 (38) Sieiro, C; Anal Quim Ser A 1984, V80, P328 HCA
 (39) Steimecke, G; Phosphorus Sulfur 1979, V7, P49 HCA
 (40) Stewart, J; The XRAY80 System 1980
 (41) Sukumar, K; Synthesis 1987, P837
 (42) Takahashi, K; J Chem Soc Chem Commun 1991, P821
 (43) Takahashi, K; J Chem Soc Chem Commun 1993, P1617 HCA
 (44) Terahara, A; Bull Chem Soc Jpn 1984, V57, P1760 HCA
 (45) Williams, J; Organic Superconductors 1992
 (46) Wudl, F; J Chem Soc Chem Commun 1970, P1453 HCA
 (47) Yamashita, Y; Angew Chem Int Ed Engl 1989, V28, P1052
 (48) Yoshida, Z; Tetrahedron Lett 1983, V24, P3469 HCA
 OSC.G 27 THERE ARE 27 CAPLUS RECORDS THAT CITE THIS RECORD (27
 CITINGS)

L13 ANSWER 7 OF 10 HCA COPYRIGHT 2010 ACS on STN

AN 107:64917 HCA Full-text

OREF 107:10649a,10652a

TI Decrosslinkable surgical materials

IN Hishida, Yasuto

PA Toyo Contact Lens Co., Ltd., Japan

SO Jpn. Kokai Tokkyo Koho, 8 pp.

CODEN: JKXXAF

DT Patent

LA Japanese

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	JP 62053664	A	19870309	JP 1985-192989	19850830
PRAI	JP 1985-192989		19850830		

AB Copolymer contg. a crosslinking agent with covalent bond capable to be opened and lost crosslinking effect by acid treatment have processable hardness before the treatment and air permeability, water absorbability, and flexibility after the treatment. A mixt. of lauryl methacrylate 36, 1,2-bis[2-(4-methacyloyloxymethyl-2-methyl-1,3-dioxolanyl)]ethane 64, and 2,2'-azobis(2,4-dimethylvaleronitrile) 0.1 part was heated 24 h at 35°, 24 h at 50°, and 24 h at 50-110°, processed and polished to give an article, swelled overnight in water, dipped overnight in 4 N aq. HCl, neutralized with 1% aq. Na2CO3, dipped overnight in water, and dipped in boiling salt soln. to give an article having Rockwell hardness 0.5, water content 37.8%, O(g) permeability 16.9 + 1010 cm3.cm/cm2.s.cmHg, and good transparency.

IT 109664-69-9

(acid treatment of, for prosthetic materials)


RN 109664-69-9 HCA

CN 2-Propenoic acid, 2-methyl-, 1,2-ethanediylbis[(2-methyl-1,3-dioxolane-2,4-diyl)methylene] ester, polymer with dodecyl 2-methyl-2-propenoate (9CI) (CA INDEX NAME)

CM 1

CRN 109664-68-8


CMF C20 H30 O8



CM 2

CRN 142-90-5

CMF C16 H30 O2



IT 109664-69-9

(acid treatment of, for prosthetic materials)

L13 ANSWER 8 OF 10 HCA COPYRIGHT 2010 ACS on STN

AN 80:37494 HCA [Full-text](#)

OREF 80:6165a,6168a

TI Synthesis of a hybrid spiro-ladder polymer

AU Bailey, William J.; Feinberg, Jay H.

CS Dep. Chem., Univ. Maryland, College Park, MD, USA

SO Polymer Preprints (American Chemical Society, Division of Polymer Chemistry) (1972), 13(1), 287-91

CODEN: ACPPAY; ISSN: 0032-3934

DT Journal

LA English

AB The dispiro bisdiene 3,4,11,12-tetramethylene-1,6,9,14-tetraoxadispiro[6.2.6.2]octadecane (I) was prepd. by heating 3,4,11,12-di(2'-dioxothiopropano)-1,6,9,14-tetraoxadispiro[6.2.6.2]- 3,11-octadecadiene [obtained from 3,4-di(hydroxymethyl)-2,5-dihydrothiophene 1,4-dioxide and 1,4-cyclohexanedione] at 200-15.deg. for 45 min. to release SO2. The I was treated with the 20-membered tetramethylene glycol maleate cyclic dimer in DMF 1 day at 5.deg., 8 days at room temp., and 1 day at 76.deg. to yield 52% polymer (II) [50601-59-7]. II was infusible but sol. in hexafluoroisopropanol at room temp. and in glutaronitrile (III) at 180.deg.. When prepd. in a III-CH2Cl2 mixt. at 5.deg. for 1 day, room temp. for 4

days, and 110.deg. for 1 day, the II was cryst. and was sol. in CH₂Cl₂ at room temp.


IT 50979-25-4P

(prepn. of)


RN 50979-25-4 HCA

CN Poly(6,6a,7,9,10,11,12,14a,15,16,21,21a,22,24,25,26,27,29,29a,30-eicosahydro-7,14,22,29-tetraoxo-1H,5H,14H,20H-[1,6,11,16]tetraoxacycloeicosino[3,4-h:13,14-h']bis[2,4]benzodioxepin-3,18-diylidene-18,18-di-1,2-ethanediyl) (9CI) (CA INDEX NAME)

PAGE 1-A



PAGE 1-B



IT 50979-25-4P

(prepn. of)

L13 ANSWER 9 OF 10 HCA COPYRIGHT 2010 ACS on STN

AN 78:16492 HCA [Full-text](#)

OREF 78:2627a,2628a

TI New spiro polymers containing five-, six-, seven-, and eight-membered cyclic ketals

AU Bailey, William John; Beam, Carles F., Jr.; Haddad, Ibrahim

CS Dep. Chem., Univ. Maryland, College Park, MD, USA

SO Polymer Preprints (American Chemical Society, Division of Polymer Chemistry) (1971), 12(1), 169-76

CODEN: ACPPAY; ISSN: 0032-3934

DT Journal

LA English


AB Tetrols prepd. were treated with cyclic diketones to give polyspiroketal ladder polymers. Spiro monomers prepd. include 7,14,21,26-tetraoxatetraspiro[5.2.2.2.5.2.2.2]hexacosane [37571-44-1], 7,14,21,25-tetraoxatetraspiro[5.2.2.2.5.2.1.2]-23-pentacosanone [37571-45-2], and cis,cis-2,3:10,11-bis(tetramethylene)-1,4,9,12-tetraoxadispiro[4.2.4.2]tetradecane [37571-46-3]. Also prepd. were 1,4-cyclohexanedione-1,1,4,4-tetrakis(hydroxymethyl)cyclohexane copolymer (I) [37450-54-7], 1,4-cyclohexanedione-1,2,4,5-tetrakis(hydroxymethyl)cyclohexane copolymer (II) [37450-55-8], 1,4-cyclohexanedione-2,2,5,5-tetrakis(hydroxymethyl)cyclopentanone copolymer (III) [37450-56-9], meso diastereomeric cis,cis-1,2,4,5-cyclohexanetetrol-1,4-cyclohexanedione copolymer (IV) [37450-57-0], 2,7-decalindione-1,2,4,5-tetrakis(hydroxymethyl)cyclohexane copolymer (V) [37450-58-1], and 1,4,5,8-tetrakis(hydroxymethyl)-1,2,3,4-tetrahydronaphthalene-1,4-cyclohexanedione copolymer (VI) [37450-59-2].

IT 39723-72-3P 39723-73-4P 39723-74-5P
39861-81-9P

(prepn. of)


RN 39723-72-3 HCA

CN Poly[(hexahydrobenzo[1,2-d:4,5-d']bis[1,3]dioxole-2,6-diylidene)-6,6-di-1,2-ethanediyl] (9CI) (CA INDEX NAME)




RN 39723-73-4 HCA

CN Poly(octahydro-1H,5H-benzo[1,2-e:4,5-e']bis[1,3]dioxepin-3,9-diylidene-9,9-di-1,2-ethanediyl) (9CI) (CA INDEX NAME)




RN 39723-74-5 HCA

CN Poly[(7-oxo-2,4,10,12-tetraoxadispiro[5.1.5.3]hexadecane-3,11-diylidene)-11,11-di-1,2-ethanediyl] (9CI) (CA INDEX NAME)



RN 39861-81-9 HCA
 CN Poly[(1,5,5a,6,7,7a,8,12-octahydronaphtho[1,8-ef:4,5-e'f']bis[1,3]dioxocin-3,10-diylidene)-10,10-di-1,2-ethanediyl] (9CI)
 (CA INDEX NAME)



IT 39723-72-3P 39723-73-4P 39723-74-5P
 39861-81-9P
 (prepn. of)

OSC.G 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD (2 CITINGS)

L13 ANSWER 10 OF 10 HCA COPYRIGHT 2010 ACS on STN

AN 77:20050 HCA Full-text

OREF 77:3365a,3366a

TI Synthesis of sulfur-containing spiro polymers

AU Bailey, William J.; Hinrichs, Robert L.

CS Dep. Chem., Univ. Maryland, College Park, MD, USA

SO Polymer Preprints (American Chemical Society, Division of Polymer Chemistry) (1970), 11(2), 598-605

CODEN: ACPPAY; ISSN: 0032-3934

DT Journal

LA English

AB C(CH₂SH)₄ (I) and cyclohexanone (using p-toluenesulfonic acid catalyst) gave 7,11,18,21-tetrathiatrispiro[5.2.2.5.2.2]heneicosane (II) [183-11-9], while 1,3-propanedithiol and 1,4-cyclohexanedione (III) gave 2,5,10,14-tetrathiadispiro[5.2.5.2]hexadecanes (IV) [34982-16-6]. I and III gave poly[3,3-bis(thiamethyl)-1,5- dithiaspiro[5.5]undecan-9-xylydene] (V) [34937-23-0], while I and 1,10-cyclooctadecanedione gave poly[3,3-

bis(thiamethyl)-1,5-dithiospiro[5.17]tricosan-15-ylidene] (VI) [34978-50-2].


II, IV, V, and VI were oxidized to their sulfones and(or) sulfoxides.

IT 34937-23-0P 36812-74-5P 36813-57-7P

(prepn. of)


RN 34937-23-0 HCA

CN Poly(2,4,8,10-tetrathiaspiro[5.5]undecane-3,9-diylidene-9,9-di-1,2-ethanediyl) (9CI) (CA INDEX NAME)




RN 36812-74-5 HCA

CN Poly[(2,4,8,10-tetraoxido-2,4,8,10-tetrathiaspiro[5.5]undecane-3,9-diylidene)-9,9-di-1,2-ethanediyl] (9CI) (CA INDEX NAME)



RN 36813-57-7 HCA

CN Poly[(2,2,4,4,8,8,10,10-octaoxido-2,4,8,10-tetrathiaspiro[5.5]undecane-3,9-diylidene)-9,9-di-1,2-ethanediyl] (9CI) (CA INDEX NAME)



IT 34937-23-0P 36812-74-5P 36813-57-7P

(prepn. of)